

Polarons in the Three-Band Peierls-Hubbard Model: An Exact Diagonalization Study

A.Dobry*

Groupe de Physique Statistique, Université de Cergy Pontoise

47 rue des Genottes, 95806 Cergy Pontoise Cedex

France

A.Greco*

Max-Planck Institut für Festkörpersforschung

Heisenbergstr. 1, D-7000 Stuttgart 80

Germany

J.Lorenzana

Laboratory of Applied and Solid State Physics. University of Groningen

Nijenborgh 4, 9747 AG Groningen

The Netherlands

J.Riera

Physics Division, Bldg. 6003, Oak Ridge National Laboratory

Oak Ridge, TN 37831

We have studied the three-band Peierls-Hubbard model describing the Cu-O layers in high- T_c superconductors by using Lanczos diagonalization and assuming infinite mass for the ions. When the system is doped with one hole, and for λ (the electron-lattice coupling strength) greater than a critical value, we found that the oxygens around one Cu contract and the hole self-traps forming a lattice and elec-

tronic small polaron. The self-trapped hole forms a local singlet analogous to the Zhang-Rice singlet in the undeformed lattice. We also studied the single-particle spectral function and the optical conductivity. We have found that the spectral weight, in general, is similar to that found in previous studies where the coupling with the lattice was absent. There is an anomalous transfer of spectral weight but, contrary to those studies, the spectral weight goes to these localized polaronic states. However this polaronic shift does not seem enough by itself to explain pinning of the chemical potential observed in real materials. The peaks in the optical conductivity are also shifted, according to the polaronic shift, in the single-particle spectral functions. We compare our results to those obtained in inhomogeneous Hartree-Fock calculations and we discuss their relation with experiments.

I. INTRODUCTION

Since the discovery of high- T_c superconductors [1], a considerable effort has been made to determine the nature of hole-doped states in the copper-oxide planes which are present in these materials. Many theoretical discussions have emphasized the importance of strong electronic correlations within these planes, leading to the study of purely electronic two-dimensional models like the one-band Hubbard model [2], the $t-J$ model [3] and three-band Hubbard models [4]. In spite of the fact that the studies on these purely electronic models are still inconclusive, there are clear experimental indications that, at least in the dilute regime, the coupling to the lattice should be taken into account [5].

The nature of doping states with and without coupling to the lattice has been studied in inhomogeneous Hartree-Fock (HF) calculations for one-band and multi-band Hubbard models [6–13]. In the three-band Hubbard model, with no coupling to the lattice, a doped hole localizes mainly in the four surrounding O sites of a Cu, nearly canceling the spin density at the central Cu [9–13]. This charge and magnetic polaron is a manifestation, at the HF level, of the singlet bound state that Zhang and Rice [3] found in their derivation of the $t-J$ model (as an effective Hamiltonian of the strongly interacting three-band Hubbard model). By including the coupling with the lattice [9,13] it was found that the O's around the central Cu forming the electronic polaron relax with a symmetric pattern [Fig. 1(b)]. Similar ideas were proposed earlier in Ref. [14]. The infrared (IR) spectra of this state [9,13], for moderate electron-lattice coupling, calculated in the random phase approximation (RPA), shows a doped-induced bleaching of the in-plane Cu-O stretching mode and a side phonon band at lower frequencies. The nature of the charge carriers and their interactions with the lattice has been probed by infrared optical absorption in photodoped [15–17] and chemically doped [18] samples. Both kind of experiments show in La_2CuO_4 , among other materials, a doping-induced bleaching of phonon modes and, in general, an intensity shift to lower frequencies. Of particular interest in La_2CuO_4 is the bleaching of the 708 cm^{-1}

Cu-O stretching mode and the appearance of a side band at 640 cm^{-1} [15,17]. Both are clearly resolved in the photodoped spectra and consistent with the data of chemically doped samples [18]. This behavior agrees well with the HF plus RPA calculations and provides a strong indication of the existence of the electronic and lattice polarons.

In spite of this success of HF plus RPA calculations, the above studies present various shortcomings. Since the electronic polaron is already self-trapped *without* coupling to the lattice, a negligible small electron lattice coupling strength λ (defined below) suffices to produce lattice relaxation. In other words, there is no competition between the delocalization energy and the self-trapping effects in this type of calculations. Another serious shortcoming comes from the fact that the Zhang-Rice wave function can not be represented by a single Slater determinant, and (at least at the HF level) a relevant part of the physics is missing. In this sense, it is important to study the interplay between electronic correlations and lattice defects or deformations without the above limitations. The electronic correlations can stabilize the lattice deformations as a consequence of the pre-existing enhancement of the effective mass. This effect has been observed in an exact diagonalization study of the Holstein-Hubbard model [19]. On the other hand, the lattice deformations can introduce anharmonic effects, further enhancing the effective mass and changing the properties of carriers with respect to those found in the framework of purely electronic models.

In this paper we study, by means of Lanczos diagonalization, the formation and properties of polarons in the two dimensional(2D) three-band Peierls-Hubbard model, which we believe is a realistic microscopic model of the Cu-O planes. The model is presented in Section II. We take the mass of the O ions equal to infinity (adiabatic approximation) for simplicity. This method is free from the above mentioned limitations of HF approximations and takes into account all the advantages of exact diagonalization studies of the three-band Hubbard models [20,21] which have reproduced fairly well the main expected spectral signatures. We discuss finite size effects in the conclusions (Section VII). By minimizing the total energy, we study the range of electron-lattice coupling at which the various lattice deformations become stable.

For any finite electron lattice coupling strength λ below a critical value (λ_c), we find that the system forms a charge density wave (CDW) with a breathing extended lattice distortion (BED) [Fig. 1(a)]. A similar distortion was found *above* a critical value in Monte Carlo simulations [22] of the Hubbard model. In Section III we discuss the phase diagram in the parameters space and give general arguments of why a BED should be found for any finite λ .

Above λ_c a polaron is formed. In what follows, we use the term polaron to designate the composite object made by the lattice deformation and the self-trapped carrier. The polaron results from static oxygen displacements as shown in Fig. 1 (b). This is in accordance with the HF picture, but in exact diagonalization calculations, the polaron appears only above λ_c whereas, as explained before, in HF it appears for any finite λ .

The polaron and the CDW are characterized by studying static magnetic and density-density correlation functions (Section IV). The presence of the self-trapped carrier manifests also in the single-particle spectral function. The evolution of the states formed by these self-trapped carriers as a function of λ is studied in Section V. Special attention is given to the issue of anomalous transfer of spectral weight and evolution of the chemical potential with doping [23,24], in view of the controversial experimental situation [25]. Finally, the effect of the self-trapped states on the optical conductivity is analyzed in Section VI. Consistently with the features of the single-particle spectra discussed above, we find a shift of the peaks to higher energies.

II. THE MODEL, PARAMETERS AND CLUSTER STUDIED

We consider the following Hamiltonian for the Cu-O planes:

$$H = \sum_{i \neq j, \sigma} t_{ij}(\{u_k\}) c_{i,\sigma}^\dagger c_{j,\sigma} + \sum_{i,\sigma} e_i c_{i,\sigma}^\dagger c_{i,\sigma} \\ + \sum_i U_i n_{i,\uparrow} n_{i,\downarrow} + \sum_{i \neq j} U_{ij} n_i n_j$$

$$+ \sum_j \frac{1}{2} K u_j^2 \quad (1)$$

Here, $c_{i,\sigma}^\dagger$ creates a hole with spin σ at the site i in the Cu $d_{x^2-y^2}$ or the O $p_{x,y}$ orbitals, $n_{i,\sigma} = c_{i,\sigma}^\dagger c_{i,\sigma}$ and $n_i = (n_{i,\uparrow} + n_{i,\downarrow})$. We have included on-site repulsions $U_i = U_d, U_p$ for holes on Cu and O atoms and a nearest-neighbor Cu-O repulsion $U_{ij} = U_{pd}$. $e_i = e_d, e_p$ are the site energies of the Cu and O atoms.

The static lattice deformation and the electron-lattice coupling are considered in the following way. Each of the patterns of deformation considered here (see next section) are defined by a set of displacements u_j of the oxygen ions along the Cu-O bonds. We assume that the nearest-neighbor Cu-O hoppings are modified by the O- ion displacements as $t_{ij} = t_{pd} \pm \alpha u_j$, where the $+$ ($-$) applies if the bond shrinks (stretches) with positive u_j . In order to study the stability of the system with respect to a given deformation of the lattice, we included the elastic energy in the Hamiltonian (last term of Eq. 1). For a given value of λ , and for a set of displacements u_j , we diagonalize exactly the electronic part of the Hamiltonian using the Lanczos algorithm. Then the total energy is minimized by varying $|u_j|$ constrained to the specific pattern.

As a reference parameter set we take $\Delta = e_p - e_d = 3, U_d = 8, U_p = 3, U_{pd} = 1, t_{pd} = 1$. For simplicity the O-O hopping t_{pp} is taken equal to zero. We take infinite masses for the O ions, so that their kinetic energy drops out. Following the estimates made in Ref. [9,13] we take $K = 32t_{pd}^2 \text{\AA}^{-2}$. In terms of the above parameters, we have defined the electron-lattice coupling as $\lambda = \frac{\alpha^2}{K t_{pd}}$.

The exact diagonalization study was made on the Cu_4O_8 cluster (12-sites) with periodic boundary conditions. We consider two cases, the half-filled system (one hole per Cu-O₂ cell), and the system doped with one extra hole.

III. STUDY OF THE STABILITY OF VARIOUS LATTICE DEFORMATIONS

As a first step in the analysis of the electron-lattice coupling, in this section we study the stability of an homogeneous distortion of the lattice (BED) [Fig. 1(a)] and the stability of a local contraction of four oxygens toward one Cu (breathing “localized” deformation or BLD) [Fig. 1(b)]. We shall also refer to this localized deformation as lattice polaron.

In the half-filling condition the lattice remains stable upon these breathing deformations, for all physically reasonable values of λ ($\lambda \lesssim 7$). A similar result was found in a Monte-Carlo study of the analogous one-band model [22] and in the HF calculations [9,13].

In general, the system can accept a lattice deformation if it gains enough local covalency to compensate the cost of elastic energy, delocalization energy, magnetic energy or Coulomb repulsion. At half-filling, the delocalization energy is obviously not relevant. We should analyze only the magnetic energy, Coulomb repulsion and covalency. Without deformation, the spin density at each Cu site is uniform and equal to a fixed value, say m . When the oxygens are displaced, the hopping integrals are modified and it is likely for charges to spend more time around the Cu sites of one sublattice and less time in the Cu sites of the other sublattice. Thus, for both Cu sites, the spin density m is reduced because, in one case, there is an increased tendency to double occupation, and in the other, the sites are becoming increasingly empty. The main increase in the energy (of order U_d) comes from the double occupancy and a large value of λ is necessary to compensate it. More complicated patterns of deformation, not studied here, can gain some covalency without increasing the double occupancy too much [12,9]. and, presumably, can destabilize the lattice for smaller λ .

The situation changes upon doping. With one doped hole, the BED is stable for every nonzero λ smaller than a certain critical value λ_c ($\lambda_c \approx 1$ for our set of parameters). The fact that the BED is stable for arbitrary small λ can be understood as follows : the ground state of the undeformed system is four-fold degenerated with momentum $k = (\pi, 0)$ or $k = (0, \pi)$ and $S_z = \pm 1/2$. Consider now the non-interacting case ($U_p = U_d = U_{pd} = 0$) with the same number of particles. The ground state is formed by five-particle Slater determinants which, for fixed S_z , can be labeled by the total momentum as, $|(\pi, 0) \rangle$ or $|(0, \pi) \rangle$. We can

construct two states $|+ \rangle = \frac{1}{\sqrt{2}}(|(\pi, 0) \rangle + |(0, \pi) \rangle)$ and $|-\rangle = \frac{1}{\sqrt{2}}(|(\pi, 0) \rangle - |(0, \pi) \rangle)$. Taking the x coordinate in the direction of the Cu₁-Cu₂ bond in Fig. 1 it is easy to see that $|+ \rangle$ will have more charge in sites 1 and 4 and $|-\rangle$ in sites 2 and 3. Clearly the BED distortion favors one of these configurations at the expense of the other. In other words, the BED lowers the symmetry of the lattice lifting the degeneracy. This conclusion is valid for the interacting case because the ground state in this system has the same symmetry and quantum numbers as in the noninteracting one.

For one doped hole, we show in Fig. 2 the ground state energy, measured with respect to the energy of the undistorted lattice, as a function of λ . Both localized and extended deformations are shown.

For $\lambda > \lambda_c$, the BLD becomes stable. The loss in elastic, Coulomb and delocalization energies is compensated by the gain in local covalency. In order to understand this crossover, we show in Fig. 3, the equilibrium lattice displacements of the oxygens along the Cu-Cu axis as a function of λ for both types of deformations. In the region $\lambda < \lambda_c$ the displacements of the ions are very small and, consequently, there are little effects of the deformation on the ground state properties. When λ becomes larger than λ_c the four displaced oxygens of the BLD undergo a sudden contraction toward their central copper. At this point this localized deformation becomes stable and the extra hole is trapped in this Cu-O₄ cluster, forming a small polaron state.

We also studied the variation of λ_c when U_d is moved away from the reference parameter set (see Fig. 4). It can be seen that λ_c increases as a function of U_d roughly linearly in the studied range ($6 \leq U_d \leq 20$). This is because BLD becomes stable by localizing the added hole (see next section) which, in turn, leads to an increasing double occupancy of the central Cu site costing an energy of order U_d . Thus, it is necessary a larger electron- lattice coupling to overcome this Coulomb repulsion.

It is worth noting that, in the strong coupling limit, this modification of the hopping integrals due to deformation leads also to a local reduction in the effective antiferromagnetic coupling J . In fact, in a perturbative calculation, J results proportional to the fourth power

of the hopping integral t and then, upon deformation $J \approx (t^2 - (\alpha u)^2)^2 < t^4$. The loss in magnetic energy, due to the lattice deformation discussed above, could also be related to this reduction in J .

IV. STATIC CORRELATION FUNCTIONS

The formation of the polaron when one hole is added to the system can be determined by analyzing, the spin and charge densities.

In Fig. 5 we show the sum of the occupancies $\langle n_i \rangle$ in the central Cu (Cu_1) and its four surrounding O involved in the BLD, as a function of λ . The total occupancy of this Cu-O₄ cluster is ≈ 2 for $\lambda > 1$ indicating the trapping of the extra hole. We also show the occupancy of the Cu_4 site, the most distant Cu from Cu_1 , and its four surrounding O.

In Fig. 6 we display $\langle S_z^2 \rangle$ in Cu_1 , Cu_4 , and in the two remaining Cu sites (Cu_2 and Cu_3), as a function of λ . The reduction of the magnetic moment at the central Cu results from the fact that the two holes, with opposite spins, are jumping back and forth between the central copper and its four surrounding oxygens. As a consequence of this there is an increased tendency to double occupancy in the central Cu. The reduction of the magnetic moment at the central Cu was also, found with qualitatively similar behavior, in Ref. [9,13].

The coupling with the lattice takes advantage of a preexisting enhanced effective mass, and produces the self-trapping of a Zhang-Rice singlet. To follow the evolution of this singlet we define the following operator: $Z_i^\dagger = (P_{i,\uparrow}^\dagger d_{i,\downarrow}^\dagger - P_{i,\downarrow}^\dagger d_{i,\uparrow}^\dagger)$ where $P_{i,\sigma}^\dagger$ creates a hole with spin σ in a symmetric combination of the four oxygen orbitals. In Fig. 7 we show the expectation value of $\langle Z^\dagger Z \rangle$ in the ground state as a function of λ . This quantity is related to the probability of finding a ZR state in the cell. We can see a rapid increases of this value when the polaron is formed. However, the double occupancy also increases at the central Cu. The amount of double occupancy could be measured from its order parameter $\langle n_{i,\uparrow} n_{i,\downarrow} \rangle$ computed at Cu_1 . We found 25% of double occupancy when the polaron is formed, while in

the undeformed lattice this value is only 6%. From this, we conclude that the polaron is a ZR singlet but of a more general character than the one found in the undeformed lattice.

V. SINGLE PARTICLE SPECTRAL FUNCTION

In this Section, we analyze the evolution of the spectral weight for single-particle excitations when the coupling to the lattice is switched on. We first review the previous known results [20,21,26,27], for $\lambda = 0$. There has been some theoretical and experimental controversy about how the Fermi level and the spectral weight evolve with doping in real materials and therefore we pay special attention to these issues, in the light of the present results. In particular, some experiments [28] suggest that the chemical potential remains pinned in the gap of the insulator as the system is doped, whereas others [25], suggest that it shifts with doping as in an ordinary semiconductor. In both cases the gap tends to fill in with transferred spectral weight.

The single-particle spectral function is defined as,

$$g_{i\sigma}(\omega) = g_{i\sigma}^>(\omega) + g_{i\sigma}^<(\omega) \quad (2)$$

where

$$g_{i\sigma}^>(\omega) = \sum_n | \langle \psi_n(N+1) | c_{i\sigma}^\dagger | \psi_0(N) \rangle |^2 \times \delta(\omega - E_0(N) + E_n(N+1)). \quad (3)$$

To obtain $g_{i\sigma}^<(\omega)$ we have to change $c_{i\sigma}^\dagger$ and $N+1$ by $c_{i\sigma}$ and $N-1$ respectively. $|\psi_n(N)\rangle$ is an eigenstate having eigenvalue $E_n(N)$ where N is the number of holes; $E_0(N)$ is the ground state energy. All the energies and gaps given below are in units of t . $g^>$ and $g^<$ describe the photoemission (PES) and inverse photoemission (IPES) spectra respectively.

Following Refs. [20] and [21], the expression for $g_{i\sigma}(\omega)$ can be evaluated numerically using the Lanczos algorithm and the continued fraction expansion formalism [29].

In Fig. 8 we show the single-particle spectral function ($g_i = g_{i\uparrow} + g_{i\downarrow}$) at the Cu and O sites in the undoped (half-filling) case $\lambda = 0$, i.e. for the undeformed lattice. The main features of this spectra have been identified in a previous study [20]. First of all, we can distinguish the lower (LHB) and upper (UHB) Hubbard bands, which are well separated by U_d plus a typical hybridization energy. Below the UHB we find the nonbonding oxygen band (NOB) with a small weight on the Cu sites. The energy difference between these two bands is slightly larger than $\Delta + 2U_{pd}$. The quantity $\Delta + 2U_{pd}$ is the (bare) charge-transfer gap expected from a ionic limit.

Between the UHB and the NOB other structure emerges. We have indicated the momentum of the states close to the Fermi energy. All of them have total spin 1/2. Due to the spatial symmetry of this lattice, the peak with momentum $k = (0, 0)$ is absent [30] but it reappears if t_{pp} is switched on [26]. These highly correlated states (CS) correspond to the Zhang-Rice singlets band [20]. The gap is given by the energy difference between UHB and CS. This energy difference is ~ 3 for our parameter set.

When the system is doped with one hole and $\lambda = 0$ (Fig. 9) the Fermi level is shifted to $\mu^+ = E_0(5) - E_0(4)$ [23,24]. This is exactly the position of the $k = (\pi, 0)$ peak in Fig. 8. This can be contrasted with the HF mean field calculations [11] where the stoichiometric system does not show spectral weight in the place where the chemical potential of the one hole doped systems sits. This is worth emphasizing because it shows that the HF results cannot be automatically invoked as an explanation of the photoemission experiments showing pinning of the chemical potential [28]. The proximity of the next available state in Fig. 9 indicates that the system is metallic. The low energy spectral weight (LESW) between the Fermi level and the gap in Fig. 9, as defined in Ref. [23,24], is equal to 1.85. This quantity should be equal to 1 for an ordinary semiconductor, and equal to 2 for the Hubbard model in the atomic limit. The value close to 2 is in agreement with the results of Ref. [23,24] and it is due to strong covalency. It is worth noting that, at the HF level, departures from an integer value of LESW (in this case 2) are not allowed. Transitions from these states to scattering states above the Fermi level were identified [11] as the origin of the mid infrared absorption

in the optical conductivity.

For $0 < \lambda < \lambda_c$, as discussed in the previous sections, the symmetry of the lattice is lowered and hence the charge tends to accumulate around the Cu_1 and Cu_4 sites (Fig. 1). This produces changes in the local g 's, but the total spectral weight remains similar to the undistorted one and the system stays metallic.

As we have shown in Section III, the BLD becomes stable in the region $\lambda > \lambda_c$. In this case, the translational invariance is broken further. In Fig. 10, we show the spectral weight for $\lambda = 1.2$. The self-trapping of the carrier around Cu_1 makes the local hole-occupied g to increase. The same quantity is larger in Cu_4 than in Cu_2 indicating that there is still some amplitude for the CS state to hop out and in from the Cu_1 , mainly to the next nearest neighbor's cell. The local g 's for the oxygens (not shown) around Cu_1 (Cu_4) increase the hole-occupied (-unoccupied) spectral weight close to the Fermi level. Apart for this feature, they are similar to the total oxygen g [Fig. 10(e)].

The levels located slightly above the Fermi level in Fig. 10 show a polaronic shift. This makes the μ^+ to sit, not in the position of the first peak in Fig. 8, like in the $\lambda = 0$ case, but somewhere in the gap. This is a direct consequence of the fact that we took an infinite mass for the O ions. If the mass were finite we expect Fig. 8 to be λ -dependent. Accordingly, some intensity will show up at the position of the new μ^+ corresponding to the polaronic configurations. However, for an infinite mass, such configurations can not be reached quantum mechanically by adding one particle to the half-filled case. Thus, their spectral weight vanish. It is tempting to associate this behavior with the experimental observation of growing spectral intensity and pinning of the chemical potential in the gap upon doping [28]. The small spectral intensity in the undoped case due to a finite O mass will be hardly measurable and hence interpreted as the gap. Nevertheless the polaronic shift close to λ_c is too small by itself to explain the effect. It can help to this interpretation when combined with other effects like impurities. The low energy spectral weight in this case is 2.06. This increase is expected from a higher covalency but, in our case, contrary to Ref. [23,24], this effect is a local one. For even larger values of λ the polaronic shift increases

further in qualitative agreement with Ref. [9,13].

VI. OPTICAL ABSORPTION

In this Section, we report our results for the optical conductivity. The calculation of this quantity in the framework of exact diagonalization follows the procedure explained in detail in Refs. [20,29] The optical absorption $\sigma'(\omega)$ or the real part of the complex conductivity $\sigma(\omega)$ is defined as:

$$\sigma(\omega) = \frac{1}{i\omega} \left[\frac{\tau^2}{N_{sites}} \langle 0 | T_y | 0 \rangle - \chi(\omega) \right] \quad (4)$$

where

$$\chi(\omega) = \frac{1}{N_{site}} \sum_n |\langle 0 | j_p | n \rangle|^2 \left[\frac{1}{\omega - (E_n - E_0) + i\eta} - \frac{1}{\omega + (E_n - E_0) + i\eta} \right]. \quad (5)$$

The quantities:

$$j_p = i\tau \sum_{i\sigma} t_{ii+y}(\{u_k\}) \left(c_{i\sigma}^\dagger c_{i+y\tau,\sigma} - c_{i+y\sigma}^\dagger c_{i\tau,\sigma} \right) \quad (6)$$

and

$$T_y = \sum_{i\sigma} t_{ii+y}(\{u_k\}) \left(c_{i\sigma}^\dagger c_{i+y\tau,\sigma} + c_{i+y\sigma}^\dagger c_{i\tau,\sigma} \right), \quad (7)$$

are the current and kinetic energy in the direction of the applied field, respectively. τ is the distance between two nearest-neighbors sites, in units of the lattice constant a .

In Fig. 11, we show the real part of the conductivity in the undoped case for $\lambda = 0$ (undeformed lattice). We can see a dominant feature formed by the peaks near $\omega = 5$. These peaks correspond to a charge transfer of a hole from Cu to O (UHB to NBO transition). As one would expect, this absorption feature appears at $\Delta + U_{pd}$ modified by a typical hybridization energy.

As we have discussed in the previous section, the minimum single particle gap (between UHB and CS) is ~ 3 . These transitions are forbidden in our cluster.

In the one-hole-doped system, and for $\lambda = 0$, it is known that there is a shift of the Fermi energy towards the CS band. Consequently, we should see a peak at low energy. Indeed, this low energy structure may be observed in Fig. 12(full line) where the first peak at $\omega \approx 2$ corresponds to transitions from CS to NBO. The structure around $\omega \approx 5$ corresponds to UHB $\rightarrow$ NBO transitions. The upper peak at ≈ 9 corresponds to CS $\rightarrow$ LHB transitions.

Now, we examine the optical conductivity in the region where the BLD is stable. In Fig. 12 (dotted line), we show the optical conductivity in the case of $\lambda = 1.2$. The most important change with respect to the undeformed case is a shift of the peaks to higher energies. Taking into account the previous analysis, we can associate this change to the polaronic shift of the single-particle levels. We also can see the set of peaks related to the UHB $\rightarrow$ NBO transitions. These peaks are approximately in the same place as before. Another shift to higher energies can be seen in the peak associated with the CS $\rightarrow$ LHB transitions. This peak appears at higher energies because it occurs between the self-trapped state and LHB.

VII. SUMMARY AND CONCLUSIONS

In this work, we have examined the self-trapping of an added hole in the Peierls-Hubbard Model using exact diagonalization in small clusters. In the presence of deformation, the hole forms a local singlet analogous to the Zhang-Rice singlet in the undeformed lattice but of a more general character, since double occupancy in Cu sites is not longer negligible.

Our calculation is complementary to inhomogeneous HF studies [9] and, in general, it confirms the qualitative behavior found in this approximation. However, there are important differences between the results obtained by these two approaches. Here we discuss these differences as well as the connection with other results in the literature.

Firstly, there is a reduction of the magnetic moments of the Cu where the doped hole is self-trapped. However, this effect is much less dramatic than in the HF calculations.

Secondly, the chemical potential for zero λ shifts with doping to a place where spectral weight already exists in the undoped compound [23,24] in contrast to the HF results. This is not the case for non-zero λ . Such an anomalous behavior arises from the lack of lattice dynamics in both our calculation and in the HF approach. The observed polaronic shift does not seem strong enough by itself to explain the observed pinning of the chemical potential.

We have shown that the spectral weight behaves in a similar manner as reported in previous studies, where the coupling with the lattice was absent [23,24]. There is an anomalous transfer of spectral weight but, contrary to those studies, this transfer is related here to local effects, i.e. it manifests in local spectral functions.

The peaks in the optical conductivity are also shifted consistently with the polaronic shift of the single-particle spectral functions. This behaviour is qualitatively similar to the HF plus RPA calculations [13] at electronic energies in *clusters of the same size*.

Thirdly, the small polaron appears in this study only above a critical value of electron-lattice coupling. This important difference with respect to HF calculations can be understood by the fact that these calculations neglect the translational motion of the correlated states and the associated kinetic energy. One expects that λ_c will be determined by the competition between the kinetic energy and the polaronic effects.

In this exact diagonalization study we have considered a quite small cluster. Unfortunately a determination of the size dependence is out of the present computational possibilities. However, if we assume that λ_c scales with the bandwidth of the Zhang-Rice states, a strong size-dependence is expected since the effective bandwidth of strongly correlated systems, for small clusters, decreases with increasing size [27].

The physics close to λ_c is highly nontrivial for a finite value of the ion mass [31]. The transition is no longer sharp but washed out by quantum fluctuations, and λ_c becomes a crossover value. The renormalization of the effective mass of quasiparticles is moderate and not exponentially large like in the large λ limit. In real materials, neither λ nor λ_c are known accurately. The fact that relatively lightly doped cuprates show already high mobilities coexisting with polaronic signatures, suggests that the system is close to λ_c . A

related scenario is that the system forms a more extended polaron with a BED distortion in a limited region, or fluctuates between both, localized and extended states [9,19].

In summary, we have given a complete characterization of the three-band Peierls-Hubbard model in small clusters in the adiabatic approximation using exact diagonalization. We have examined some properties which can not be properly considered within HF calculations, which have been so far extensively used to study these systems. We believe that these effects can be generic to a wide class of materials.

Acknowledgments

A.D. and J.L. are supported in this work by postdoctoral fellowships granted by the European Economic Community. A.D. is also indebted to Fundación Antorchas of Argentina for traveling support. A.G. is grateful to A. Aligia, C. Balseiro, P. Etchegoin, I. Mazin and R.Zeyher for useful discussions. Special thanks are due to P. Horsch for a critical reading of the manuscript. A.G. is supported by a fellowship from the Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina. J.L. is in debt with G. Sawatzky for many discussions on how the chemical potential and the spectral weight evolve with doping in a strongly correlated system and with F. Barriquand for a critical reading of the manuscript. During the realization of this work, J.R. was supported by a fellowship from the Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina.

*Permanent address: Instituto de Física Rosario (CONICET-UNR) and CIUNR. Bv. 27 de Febrero 210 Bis. 2000 Rosario, Argentina.

REFERENCES

- [1] J. G. Bednorz and K. A. Muller, Z. Phys. B**46**, 189 (1986).
- [2] P.W.Anderson, Science, **235**, 1196 (1987).
- [3] F. C. Zhang and T. M. Rice, Phys. Rev. B**37**, 3759 (1988). E. Dagotto et al, Phys. Rev.B **45**, 10741 (1992).
- [4] V. Emery, Phys. Rev. Lett.**58**, 2794 (1987); J.Hirsch, Phys. Rev.Lett.**59**, 228 (1987); C.Varma et al, Solid State Commun. **62**, 681 (1987).
- [5] For a recent overview see Proceedings of the Conference *Lattice effects in High-Tc Superconductors*, Santa Fe, New Mexico, January 1992, edited by Y. Bar-Yam et al. (World Scientific, Singapore).
- [6] J. Lorenzana and L. Yu, Phys. Rev. B **43**, 11474 (1991). J. Lorenzana, PhD Thesis (International School for Advanced Studies, Trieste, 1992).
- [7] J. Lorenzana and L. Yu, Mod. Phys. Lett. B **5**, 1515 (1991).
- [8] M. Inui and P. B. Littlewood, Phys. Rev. B **44**, 4415 (1991).
- [9] K. Yonemitsu, A.R. Bishop, and J. Lorenzana, Phys. Rev. Lett. **69**, 965 (1992).
- [10] J. B. Grant and A. K. Mc Mahan, Phys. Rev.B **46**, 8440 (1992).
- [11] J. Lorenzana and L. Yu, Phys. Rev. Lett. **70**, 861 (1993).
- [12] K. Yonemitsu, A.R. Bishop, and J. Lorenzana, Phys. Rev. B **47**, 8065 (1993).
- [13] K. Yonemitsu, A.R. Bishop, and J. Lorenzana, Phys. Rev. B **47**, 12 059 (1993). Idem, Ref. [5].
- [14] M. J. Rice and Y. R. Wang, Phys. Rev. B **36**, 8794 (1987).
- [15] Y. H. Kim, C. M. Foster, and A. J. Heeger, Phys. Scr. T **27**, 19 (1989).

- [16] X. Wei, C. Chen, Z.V. Vardeny, C. Taliani, R. Zamboni, A.J. Pal, and G. Ruani, *Physica C* **162-164**, 1109 (1989).
- [17] For a recent review see, G. Yu and A. Heeger, Submitted to *Rev. Mod. Phys.* This reference discuss the remarkable similarity between the photodoped and chemically doped experiments.
- [18] G. A. Thomas, D. H. Rapkine, S-W. Cheong, and L. F. Schneemeyer, *Phys. Rev. B* **47**, 11396 (1993).
- [19] J. Zhong, et al, *Phys. Rev. Lett.* **69**, 1600 (1992).
- [20] J. Wagner, et al, *Phys. Rev. B* **43**, 10517 (1991).
- [21] P. Horsch, et al, *Physica C* **162-164**, 783 (1989).
- [22] A. Muramatsu and W. Hanke , *Phys. Rev. B* **38**, 878 (1988).
- [23] H. Eskes, M.B.J. Meinders, and G. A. Sawatzky, *Phys. Rev. Lett.* **67**, 1035 (1991).
- [24] M.B.J. Meinders, H. Eskes and G. A. Sawatzky, preprint.
- [25] M.A. van Veenendaal, R. Schlatmann, G. A. Sawatzky and W. A. Groen, *Phys. Rev. B* **47**, 446 (1993).
- [26] H. J. Schmidt and Y. Kuramoto, *Phys. Rev. B* **42**, 2562 (1990).
- [27] Y. Ohta, K. Tsutsui, W. Koshibae, T. Shimozato and S. Maekawa, Evolution of the in-gap state in High-Tc Cuprates, preprint.
- [28] J. W. Allen et al., *Phys. Rev. Lett* **64**, 595 (1990).
- [29] E. Gagliano and C.Balseiro, *Phys. Rev. B* **38**, 11761 (1988).
- [30] A. Aligia, private communication. For the spatial symmetries of the states, see C. Batista and A.A. Aligia, *Solid State Commun.* ,**83**, 419 (1992). Idem, submitted to *Phys. Rev. B*.

- [31] H. de Raedt, A. Lagendijk, Z. Phys.B **65**, 43 (1986); D. Feingberg, S. Ciuchi, F. de Pascuale, Int. J. Mod. Phys. **4**, 1317 (1990); D. Feingberg, J. Ranninger, Phys.Rev. A **30** (1984); *ibid*, **33** 3466 (1986).

FIGURES

FIG. 1. Patterns of displacements of the oxygen ions along the Cu-O bonds for: (a) Breathing Extended Deformation (BED) and (b) Breathing Local Deformation (BLD).

FIG. 2. Ground state energy as a function of λ for both, BED and BLD. The energy is measured with respect to the energy of the undistorted lattice.

FIG. 3. Equilibrium lattice displacements of the O's along the Cu-Cu axis as a function of λ for both type of deformations (BED and BLD).

FIG. 4. λ_c as a function of U_d .

FIG. 5. Occupancies of the CuO_4 cluster as a function of λ in the BLD. We show occupancies in both Cu_1 and Cu_4 clusters.

FIG. 6. $\langle S_z^2 \rangle$ as a function of λ in the BLD. We show $\langle S_z^2 \rangle$ in Cu_1 , Cu_2 and Cu_4 clusters.

FIG. 7. $\langle Z^\dagger Z \rangle$ as a function of λ in the BLD.

FIG. 8. Single particle spectral function for the half-filled case in (a) Cu and (b) O sites. The full and dashed lines correspond to PES and IPES experiments, respectively. The spectra is independent of λ since for reasonable values of λ the lattice is undistorted.

FIG. 9. Single particle spectral function in (a) Cu and (b) O sites for the 1 hole doped case and $\lambda = 0$. The full and dashed lines correspond to PES and IPES experiments, respectively.

FIG. 10. Single particle spectral function for $\lambda = 1.2$. The full and dashed lines correspond to PES and IPES experiments, respectively. We show the spectra in non equivalent Cu's (a,b,c) and the total spectra in Cu (d) and O (e).

FIG. 11. Real part of the conductivity in the undoped case for $\lambda = 0$.

FIG. 12. Real part of the conductivity in the one hole doped case for $\lambda = 0$ (full line) and for $\lambda = 1.2$ (dotted line).